



Contribution ID: 34

Type: **Contributed talk**

# Unravelling the Stratification of the Chromospheric Magnetic Field Using the $H\alpha$ Line

*Tuesday, January 21, 2025 9:10 AM (15 minutes)*

The  $H\alpha$  line is widely utilized for studying the solar chromosphere, but there is a scarcity of polarimetric studies aimed at inferring magnetic fields. One of the many reasons could be that there are no polarimetric studies of the  $H\alpha$  line utilizing 3-D radiative transfer, and earlier 1-D radiative transfer studies suggested a significant contribution of the photospheric fields in the Stokes V profiles. In our recently published work, Mathur et al. 2023, using spectropolarimetric data of a pore simultaneously recorded in the  $H\alpha$  and Ca II 8542 Å lines we investigated the potential of  $H\alpha$  Stokes V profiles in determining chromospheric magnetic fields. Our findings suggested that the line core of the  $H\alpha$  line probes the chromospheric magnetic field. However, the previous study was limited to a small pore. In this study, using spectropolarimetric observations of an active region recorded simultaneously in the  $H\alpha$  and Ca II 8662 Å lines, we inferred the stratification of the chromospheric magnetic field. The sunspot exhibits multiple structures, viz., 4 umbras and a lightbridge and a region where Ca II 8662 Å line core is in emission. The  $H\alpha$  line core image also displays brightening in the emission region, a signature of localized heating, with the spectral profiles showing elevated line cores. Consistent with the Mathur et al. 2023, we found that the magnetic field inferred from the  $H\alpha$  line core is consistently smaller than that inferred from inversions of the Ca II 8662 Å line at  $\log \tau_{500} = -4.5$ , however, in contrast with Mathur et al. 2023, uncorrelated. The field strength and morphology inferred in the heating region from the inversions at  $\log \tau_{500} = -4.5$  is comparable to that of at  $\log \tau_{500} = -1$ . There is also a good agreement with the field strengths at  $\log \tau_{500} = -1$  with that inferred from WFA over  $H\alpha$  full spectral range, except in the heating region. In addition, the fields inferred in the heating region from the WFA over  $H\alpha$  line core and full spectral range are similar in strengths and morphology. In addition, we have also performed a theoretical study of synthesizing the polarization profiles of the  $H\alpha$  line in 3D. We show that the line of sight magnetic field retrieved is sensitive to  $\log \tau_{500} = -5.7$ , which is at higher heights compared to the Ca II IR line. Thus, we suggest that the line core of the  $H\alpha$  line always probes the chromospheric magnetic field at higher heights than that probes by the Ca II IR triplet. In case of heating events, the full  $H\alpha$  line becomes sensitive to the chromospheric magnetic field instead of just the line core. Consequently, the  $H\alpha$  line spectropolarimetry is a valuable diagnostic for studying the chromosphere, especially in regions with localized heating, where the Ca II IR triplet lines probe deeper layers of the solar atmosphere.

## Contribution Type

### Theme

Solar Magnetism in High-Resolution

**Primary author:** MATHUR, Harsh (Indian Institute of Astrophysics)

**Co-authors:** JOSHI, Jayant (Indian Institute of Astrophysics); K, Nagaraju (Indian Institute of Astrophysics); Dr YADAV, Rahul (Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA)

**Presenter:** MATHUR, Harsh (Indian Institute of Astrophysics)

**Session Classification:** High Resolution Observations of Solar Magnetic Fields